

Gas Chromatographic Analysis of the Phyto-Constituents and the Assessment of the Anti-Microbial Properties of the Leave Extracts of Nigeria-Grown *Ginkgo biloba*

Ubaoji, K.I.¹, Nwosu, O.K², Agu, K.C.³, Nwozor, K.O.⁴, Ifedilichukwu, N.H.⁵, Okaka, A.N.C.⁶

¹ Department of Applied Biochemistry, Nnamdi Azikiwe University, Awka, Nigeria.

² National Biosafety Management Agency, Abuja, Nigeria

³ Department of Applied Microbiology, Nnamdi Azikiwe University, Awka, Nigeria.

⁴ Department of Biochemistry, Federal University of Gusau, Nigeria.

⁵ Department of Medical Biotechnology, National Biotechnology Development Agency, Abuja.

⁶ Department of Applied Biochemistry, Nnamdi Azikiwe University, Awka, Nigeria.

Article Info

Article history:

Received: 07 September 2020

Revised: 18 October 2020

Accepted: 22 October 2020

Keywords:

Ginkgo biloba,
Phytochemicals,
Antibacterial,
Antifungal, Gas
Chromatography,
Microorganisms.

Paper Type :

Research Article

Corresponding Author:

Nwosu, O.K

Email:

nwosuonyeka6@gmail.com

Abstract

Purpose: *Ginkgo biloba* (GB) leaves have been recognized for their therapeutic effects, dating back to traditional Chinese medicine where the ground leaves were used to treat several health problems. GB extracts are renowned for their medicinal properties in Asia, Europe and South America. However, GB leaves are rarely grown and used in Nigeria. In this study, we screened the phytochemical constituents and antimicrobial activities of GB leaves (aqueous and ethanol extracts) grown in Enugu, Enugu State, South East Nigeria.

Study Design: Experimental

Materials and Methods: Quantitative analyses of phytochemicals were done using gas chromatography while disk diffusion and broth dilution methods were used to determine the antimicrobial activities.

Results: The phytochemical analysis of aqueous and ethanol extracts showed high quantity of flavonoids (anthocyanin, rutin, epicatechin, kaemferol and catechin), alkaloids (lunamarine and ribalindine) and saponin while the moderate quantity of steroids and terpenes, tannin and phenols were also observed. The phytochemical analysis also showed very little concentration of antinutrients (phytate and oxalate) in both extracts. The ethanol extract showed some degree of antibacterial and antifungal activities as observed from their zones of inhibition (mm). Activity was observed on gram-positive bacteria; *Staphylococcus aureus* ($15.5+0.71\text{mm}$) and less activity on gram-negative bacteria; *Escherichia coli* ($10.5+1.41\text{mm}$) and *Pseudomonas aeruginosa* (no activity). Also there was an activity on yeast; *Candida albicans* ($16.5+0.71\text{mm}$) and less activity on moulds; *Penicillium cyclopium* ($9+1.41\text{mm}$) and *Aspergillus fumigatus* (no activity). The aqueous extract showed no antimicrobial activity on both bacteria and fungi.

Conclusion: This study revealed the medicinal potentials of *Ginkgo biloba* leaves grown in Nigeria. The findings, therefore, will be useful for pharmaceutical industries and medical practice in Nigeria and beyond for the provision of good health to individuals.

1. Introduction

For many years, plants have been used as medicine because they are an important source of many biologically and therapeutically active products. These plants have been

implicated in the treatment of several ailments but some of them may have undesirable effects in the body. Even though medicinal plants have provided biologically relevant products for centuries, they still serve as a source for new medicines, thereby increasing the global interest in ethnopharmacological studies (Damery et al., 2011). The increasing motivation for these ethnopharmacological studies in recent times is based on how traditional healers (herbalists) use plants therapeutically. The essence of such studies is to identify or discover interesting properties and constituents of plants that make them therapeutic or medicinal. The medicinal properties of the plants are mainly attributed to the presence of various simple and complex organic compounds in them (Raaman, 2008). These compounds are called secondary metabolites. It is due to these compounds that many medicinal plants are said to be therapeutic including being anti-microbial.

These secondary metabolites are non-essential nutrients in such that they are not required by animal for sustaining life (Raaman, 2008). Unlike primary metabolites, the absence of secondary metabolites does not result in immediate death, but in the long-term impairment of the organism's survival/fecundity or aesthetics or, perhaps, in no significant change at all (Kennedy & Whightman, 2011). The common secondary metabolites usually obtained from plants are flavonoids, alkaloids, anthraquinone, cardiac glycoside, phenols, terpenes, steroids, saponin, tannin, reducing sugar and pyrolizidine (Dillard & German, 2000).

Microbes or microorganisms are microscopic living organisms that are single-celled or multicellular. Many of these microbes are especially dangerous for the very young, frail elderly, immun-compromised or critically ill individuals, as well as patients with particular physiological or dermatological disorders (Jayaprakash et al., 2003). Notably, healthy individuals are also at risk, and this is most clearly evident in less affluent regions in the world like Nigeria, where infectious diseases are still a major cause of morbidity and mortality. Antimicrobials are all agents (including plant parts) that act against all types of microorganisms- bacterial (antibacterial), fungi (antifungal), viruses (antiviral) and protozoa (antiprotozoal). These antimicrobials according to Peach et al. (2013) have different modes of action, owing to the nature of their structure and degree of affinity to certain target sites. They include inhibitors of cell wall synthesis, inhibitors of cell membrane functions, inhibitors of protein synthesis and inhibitors of nucleic acid synthesis among others.

An ancient Chinese tree that had appeared more than 250 million years ago, and the only surviving member of Ginkgoaceae family; *Ginkgo biloba*, has been used extensively by traditional and visionary healers around the world, especially Asia and South America due to its acclaimed therapeutic functions (DeFeudis, 1998). The leaves of *Ginkgo biloba* tree have attracted attention as agents for improving circulation, particularly cerebral circulation which may lead to improved mental functions (Wadsworth & Koop, 2001). Kwon et al. (2004) reported that *Ginkgo biloba* leaf extracts have been implicated in the treatment of poor circulation, impotence, heart diseases, eye diseases, chronic cerebral insufficiency, short term memory loss (like Alzheimer's diseases), depression and dementia. This paper is designed to analyze the phytochemical constituents and determine the anti-microbial properties of Nigerian grown *Ginkgo biloba* leaves (aqueous and ethanol extract).

2. Methodology and Procedures

Collection and Extraction of Plant Leaves

Healthy fresh leaves of *Ginkgo biloba* were collected from *Ginkgo biloba* tree at Park Avenue, GRA, Enugu, Enugu State, Nigeria in a large quantity. The plant was identified and authenticated by a plant taxonomist in the Botany Department of Nnamdi Azikiwe University, Awka. The voucher specimen was deposited in the Herbarium of the Department. The leaves were dried at room temperature for three weeks. The dried leaves were later ground into a fine powder using a local steel grinder. To obtain an aqueous extract, 30g of the ground leaves was mixed with 300ml of distilled water. Then 30g of the ground leaves was mixed with 80% (v/v) ethanol to obtain an ethanol extract. Both mixtures were refluxed in a water bath at 65°C for 1 hour and filtered using Whittman filter paper No. 1 followed by the evaporation of the filtrate to slurry form using a rotary evaporator.

Phytochemical Quantification

Quantification of the phytochemical constituents in the aqueous and ethanol extracts was done using gas chromatography, fitted with flame ionization detector (GC-FID) employing Martin and Synge (1941) method and modified by Ujowundu et al. (2015).

Procedure:

For each extract, 0.5ml was introduced into separating funnels containing 0.5g of sodium sulphate. Then 20ml of N-hexane was added to each column and was left to stand for a few minutes after which the N-hexane layer was extracted and used for the GC-FID analysis.

Conditions for the Quantification:

The quantification of the phytochemicals was performed on a BUCK M910 Gas Chromatograph equipped with a flame ionization detector. A RESTEK 15 meter MXT-1 column (15 x 250 x 0.15 μ m) was used. The injector temperature was 280°C with a splitless injection of 2 μ l of the sample and a linear velocity of 30cms $^{-1}$. Helium 5.0_{pa.s} was the carrier gas with a flow rate of 40mlmin $^{-1}$. The oven operated initially at 200°C, it was heated to 330°C at a rate of 3°C min $^{-1}$ and was kept at this temperature for 5 minutes. The detector operated at a temperature of 320°C.

The phytochemicals were then determined by the ratio between the area and mass of internal standards and the area of the identified phytochemicals. The concentration of the different phytochemicals was expressed in μ g/ml.

Antimicrobial Analysis

Test Organisms:

A total of three pathogenic bacteria namely: *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and three pathogenic fungi, *Candida albicans* *Aspergillus fumigatus*,

Penicilliumcyclopium were obtained from the Reference Laboratory section of the Department of Applied Microbiology and Brewing, NnamdiAzikiwe University Awka, Anambra State, Nigeria. These organisms were maintained on Nutrient Broth for Bacteria and Sabouraud Dextrose Broth for fungi.

Antimicrobial Susceptibility Test:

This was determined by the modified method described by Agu et al. (2013). Plates that had confluent and/or semi-confluent growth were selected for the antimicrobial susceptibility tests. The disk diffusion method was used to assay the effect of the extracts on the various microorganisms. Mueller-Hinton Agar was used for the bacteria and Sabouraud Dextrose Agar (SDA) for the fungus. Twenty four hours broth cultures of the test organisms were serially diluted, then 10^{-1} and 10^{-2} dilutions were used to seed the fungal isolates, while 10^{-2} was used to seed the bacterial plates. Then 0.1ml of the appropriate dilution of the broth culture of each microorganism was uniformly spread using a sterile glass spreader on the surface of the media, and sterile filter papers were soaked in the neat *Ginkgo biloba* extracts (aqueous and ethanol) that is in slurry form and placed on two points on each petri dish. Incubation was done at room temperature for 24 hours for bacteria and 48 hours for fungi. Clear zones of inhibition around the wells indicated antimicrobial activities of the extracts against the test organisms. The diameters of the zones of inhibition were measured and recorded in millimetres. Negative controls were set up with sterile water and positive controls were set up using 0.5% Nystatin for fungi and 0.5% Ciprofloxacin for bacteria.

Determination of Minimum Inhibitory Concentration (MIC):

The Modified method of Pallotaet al. (2007) was used for this study. The Minimum Inhibitory Concentration (MIC) was determined by the broth dilution method. Different test tubes containing Nutrient Broth (bacteria) and Sabouraud Dextrose Broth (fungi), which contained 20%, 40%, 60% and 80%, of the extracts were inoculated with known amounts of the bacterial and fungal isolates, with a dilution factor of 10^{-2} for bacteria and 10^{-1} and 10^{-2} for fungi and controls (negative and positive) were set up also. The plates were incubated at room temperature for 24 hours for both bacteria and fungi in a metabolic rotary shaker (220 revs/min). Thereafter, the incubated test tubes were then sub-cultured onto sterile freshly prepared plates and incubated for 24 hours for bacteria and 48 hours for fungi. At the end of the incubation period, the plates were counted and the total microbial count recorded in CFU/ml. The plate with the lowest count was recorded as the MIC. The MIC is defined as the lowest concentration required to arrest the growth of the micro-organism at the end of 24 h or 48 h of incubation.

3. Results and Discussion

Some data obtained in this study were evaluated using the Statistical Package for Social Sciences (SPSS). The values were expressed as a mean of duplicate determination \pm standard deviation. Analysis of variance (ANOVA) was done at 0.05 level of significance, if $P < 0.05$, there is a significant difference.

Table 1: GC-FID Quantitative Analysis of Phytochemical Compositions of Aqueous Extract of *Ginkgo biloba* leaf

Components	Subclass	Concentration ($\mu\text{g}/\text{ml}$)	% Concentration	Retention time (mins)
Flavonoids				
	Anthocyanin	0.745	0.455	1.883
	Rutin	27.651	16.871	8.750
	Epicatechin	3.674	2.242	15.696
	Kaempferol	45.474	27.746	39.246
	Catechin	0.858	0.523	45.253
Total			47.837	
Alkaloids				
	Lunamarine	26.669	16.272	18.260
	Ribalnidine	1.661	1.013	31.460
Total			17.285	
Steroids and Terpenes				
	Sapogenin	19.859	12.117	29.116
Saponin		21.316	13.006	24.106
Tannin		7.986	4.873	4.766
Phenol		7.416	4.525	12.666
Phytate		0.586	0.358	32.336

Source: Authors

Table 2: GC-FID Quantitative Analysis of Phytochemical Compositions of Ethanol Extract of *Ginkgo biloba* leaf

Components	Subclass	Concentration ($\mu\text{g}/\text{ml}$)	% Concentration	Retention time (mins)
Flavonoids				
	Anthocyanin	0.746	0.369	1.883
	Rutin	27.660	13.682	8.750
	Epicatechin	3.685	1.823	15.696
	Kaempferol	45.457	22.486	39.246
	Catechin	41.482	20.519	45.350
Total			58.879	
Alkaloids				
	Lunamarine	26.775	13.244	18.260
	Ribalnidine	1.661	0.916	31.413
Total			14.160	
Steroids and Terpenes				

	Sapogenin	20.004	9.895	29.116
Saponin		21.334	10.553	24.106
Tannin		4.529	2.240	4.903
Phenol		7.430	3.675	12.666
Oxalate		0.650	0.322	4.596
Phytate		0.555	0.275	32.353

Source: Authors

Table 3 Antibacterial Susceptibility Screening of *Ginkgo biloba* leaf Extracts

Zone of Inhibition (mm)			
Isolates	Aqueous Extract	Ethanol Extract	Positive Control
<i>Staphylococcus aureus</i>	-	15.5 ± 0.71*	39 ± 1.41
<i>Escherichia coli</i>	-	10.5 ± 0.71*	26 ± 2.12
<i>Pseudomonas aeruginosa</i>	-	-	42 ± 2.83

Data are the means of duplicate determination ± standard deviation. The positive control is Ciprofloxacin. The negative control used was sterile water which showed no inhibition. *p<0.05 when compared with positive control.

Source: Authors

Table 4 Antifungal Susceptibility Screening of *Ginkgo biloba* leaf Extracts

Zone of Inhibition (mm)			
Isolates	Aqueous Extract	Ethanol Extract	Positive Control
<i>Aspergillus fumigatus</i>	-	16.5 + 0.71*	41 + 1.41
<i>Candida albicans</i>	-	-	34.5 + 2.12
<i>Penicillium cyclopium</i>	-	9+ 1.41*	36.5 + 2.12

*Data are the means of duplicate determination \pm standard deviation. The positive control is Nystatin. The negative control used was sterile water which showed no inhibition. * $p < 0.05$ when compared with positive control.*

Source: Authors

Table 5 Minimal Inhibitory Concentration (MIC) (%) of the Ethanol Extract of *Ginkgo biloba* leaf on the Test Bacteria

Isolates	Colony Count After 24 hours(mm)			
	20% Conc	40% Conc	60% Conc	80% Conc
Ethanol Extract				
	20 \pm 2.83	NC	NC	NC
<i>Staphylococcus aureus</i>				
Positive Control				
	3 \pm 0.00	NC	NC	NC
Ethanol Extract				
	16 \pm 2.83	5 \pm 0.00	2 \pm 0.00	1 \pm 0.00
<i>Escherichia coli</i>				
Positive Control				
	12 \pm 1.41	NC	NC	NC

Data are the means of duplicate determination \pm Standard deviation. The positive control is Ciprofloxacin. The negative control used was sterile water which showed no inhibition. NC = No count.

Source: Authors

Table 6: Minimal Inhibitory Concentration (MIC) (%) of the Ethanol Extract of *Ginkgo biloba* leaf on the Test Fungi

Isolates/DF	Colony Count After 48 hours (mm)			
	20% Conc	40% Conc	60% Conc	80% Conc
Ethanol Extract				
	34 \pm 1.41	9 \pm 1.41	1 \pm 0.00	NC
<i>Candida albicans/10⁴</i> Positive Control				
	7 \pm 1.41	NC	NC	
Ethanol Extract				
	15 \pm 2.83	4 \pm 0.00	NC	NC

Candida albicans/10²Positive Control

4 ± 1.41	NC	NCNC
Ethanol Extract		
14 ± 1.41	3 ± 1.41	NC
Penicilliumcyclopium/10 ⁻¹ Positive Control		
NC	NCNCNC	
Ethanol Extract		
7 ± 1.41	2 ± 0.00	NC
<i>Penicilliumcyclopium</i> /10 ⁻² Positive Control		
NC	NCNCNC	

Data are means of duplicate determination ± Standard deviation. Positive control is Nystatin. The negative control used was sterile water which showed no inhibition. DF = Dilution factor, NC = No count.

Source: Authors

Discussion

In this study, gas chromatography linked with flame ionization detector (GC-FID) showed the quantities of 12 and 13 specific phytochemicals in the aqueous and ethanol extracts respectively, as seen in tables 1 and 2. The identification of these specific phytochemicals was done by comparison of the GC- retention time of the standard phytochemicals which was saved in the data system library. Flavonoids content was observed to be in the highest quantity in both extracts considering the concentration levels of the different components of flavonoids like epicatechin, kaempferol, anthocyanin, catechin and rutin. Catechins have shown to possess antibiotic properties due to their role in disrupting a specific stage of the bacterial DNA replication process (Gradisar et al., 2007). It may have contributed immensely to the many reported antibacterial effects of the leaf. Rutin has been reported to have clinically relevant functions that include anti-inflammatory, antihaemorrhagic activity and strengthening of capillary permeability and stabilization of platelets (Guo and Wei, 2008).Kaempferol and anthocyanin are antioxidant components and could have aided in the massive reports of the antioxidant properties of the leaves. Anthocyanin has been reported to be involved in the improvement of vision, decreased platelet aggregation and neuroprotective effects (Tyagi et al., 2010). The specific alkaloid; lunamarine has shown to be immunomodulatory and anti-amoebic activity, particularly against *Entamoeba histolytica*. Alghabban (2014) in his study confirmed that the *Ginkgo biloba* leaf extract inhibited and treated *Entamoeba histolytica* infection. Tannin was also identified as shown in Table 1 and 2 for aqueous and ethanol extracts. Tannin has been reported to be an essential antioxidant, antimicrobial and anti-cancer activities (Dillard and German, 2000). Saponin content in aqueous and ethanol extracts is quite significant with concentrations of 21.3164µg/ml and 21.3336µg/ml respectively. The saponins are generally

known to exert hypoglycaemic and anticholesterol effect through intra-luminal physiochemical interactions (Price et al., 1987). Saponin has also been implicated to act as antibacterial agents (Lacaille-Dubois & Wagner, 2000).

This study as with many other studies on *Ginkgo biloba* had indicated the high content of phenols in the leaf and had contributed immensely with steroids and terpenes (sapogenin) and flavonoids to its highly recognized antioxidant effect. The study also identified very low content of anti-nutrients (oxalate and phytate) in both extracts which are very good for consumption. The study has shown that the extracts contain appreciable phytochemicals that exact many therapeutic functions, it can be said that the extracts may be safe for human consumption for medicinal purposes. However, it is pertinent to state that the percentage of each constituent in the extracts can vary according to the country where it was grown and the season in which the plant leaves were harvested (Sati et al., 2013).

Table 3 and 4 showed that the aqueous extract did not show any antibacterial and antifungal activity on the three bacteria and fungi. Ethanol may be a better solvent for the extraction of the secondary metabolites from the leaves especially catechin, as shown in comparison between table 1 and 2. Catechin may have being the leaf's constituent that easily diffused into the bacterial and fungal cell wall. The reason for the reduced effect of the extracts on gram-negative bacteria and mould may have been due to the complexity of the cell wall of gram-negative bacteria and the central core of the mould cell wall that is a branched β 1,3,1,6 glucan linked to chitin via β 1, 4 linkage (Latge, 2007) respectively. Majorly, β 1, 3 glucan synthase inhibitors is used clinically to target *Aspergillus fumigatus* and other mould fungi (Latge, 2007). This result is in consonance with Sati and Joshi (2011) who confirmed the antibacterial effect of *Ginkgo biloba* ethanol extract of Indian origin on many bacteria including *Escherichia coli* that showed low inhibition. The result is also in consonance to those of Sati et al., (2012) who confirmed that the aqueous leaf extract of *Ginkgo biloba* from India did not show any activity on gram-negative and gram-positive bacteria, and mould and yeast whereas other extracts including ethanol showed antibacterial and antifungal effect.

In table 5, the colony count was high in 20% dilution of the *Ginkgo biloba* ethanol extract for *Staphylococcus aureus* and *Escherichia coli*. This is due to the low concentration of the extract, however 40%, 60% and 80% concentrations for *Staphylococcus aureus* showed no colony count indicating that 40% dilution (0.4ml of extract in 0.6ml of broth) extract may have been able to inhibit and kill *Staphylococcus aureus*. As for the *E.coli*, 40%, 60% and 80% dilution concentration of the extract showed a decrease in the colony count in that sequential order. Therefore it can be said that at 40% dilution concentration, the extract may inhibit *E.coli* to the greatest minimum, and at 80% dilution concentration, the extract may kill *E.coli*.

In table 6, the colony count was high in 20% of 10^{-1} dilution factor with 34 ± 1.41 count and 20% of 10^{-2} dilution factor with 15 ± 2.83 count of *Candida albicans*. This is due to the low concentration of the extract while 40% and 60% concentration for 10^{-1} dilution factor; the count decreased showing greater inhibition on *Candida albicans*. The 80% concentration of the extract may be able to kill *Candida albicans*. It was the same for 10^{-2} dilution factor with 60% concentration that may be able to inhibit and kill *Candida albicans*.

The same inhibition based on concentration was observed in *Penicilliumcyclopiumas* in *Candida albicans*. However, at 60% concentration of the ethanol extract, it was able to inhibit and kill *Penicilliumcyclopium* at 10^{-1} and 10^{-2} dilution factors.

The inhibition activity of the ethanol extract of *Ginkgo biloba* leaf signifies the leaf antimicrobial effect and this might be due to presence of variety of active compounds in the extract such as tannin, saponin, alkaloids and flavonoids as suggested by Abo *et al.*, (2000). This study had established the potential of using *Ginkgo biloba* ethanol leaf extract of Nigerian origin for microbial treatment.

Conclusion and Suggestion

In conclusion, this study revealed that aqueous and ethanol leaf extracts of *Ginkgo biloba* from Nigeria possess phyto-constituents of health importance and have shown to possess antibacterial and antifungal properties. The importance of natural products with phyto-constituents of health implication and antimicrobial properties cannot be over-emphasized; therefore greater awareness should be vigorously pursued to effectively appropriate the rich chemical constituents and anti-microbial properties of *Ginkgo biloba* found in Nigeria.

Conflict of Interest

The authors of the article declare no conflict of interest.

Funding:

The authors received no funding for conducting this study.

References

- Abo, K.A, Adeyemi, A.A and Jegede, I.A. (2000). Spectrophotometric Estimation of Anthoquinone Content and Antimicrobial Potential of Extracts of some *Cassia* Species Used in Herbal Medicine in Ibadan. *Science Forum*,3(2): 57-63.
- Agu, K.C., Igweoha, C.A and Umeh, C.N. (2013).Antimicrobial Activity of the Ethanolic and Petroleum Ether Extracts of Tangerine Seed on Selected Bacteria.*International Journal of Agriculture and Bioscience*,2(1): 22-24.
- Alghabban, A.J.M (2014). *Ginkgo biloba* Leaf Extract Inhibits and Treated *Entamoeba histolytica* Infection. *Nature and Science*,12(2): 63-68.
- Damery, S., Gratus, C and Grieve, R. (2011). The Use of Herbal Medicine by People with Cancer: A Cross Sectional Survey. *British Journal on Cancer*, 104(6): 927-933.
- Defeudis, F.V. (1998). *Ginkgo biloba* Extract (EGb761), From Chemistry to Clinic.Ullstein Medical, USA
- Dillard, C.J and German, J.B. (2000). Phytochemicals: Nutraceuticals and Human Health. *Journal of Science of Food and Agriculture*,80: 1744-1756.

- Gradisar, H., Pristovsek, P., Plaper, A and Jerala, R. (2007). Green Tea Catechins Inhibit Bacterial DNA Gyrase by Interaction with its ATP Binding Site. *Journal of Medicinal Chemistry*, 50: 264-271.
- Guo, R and Wei, P. (2008). Studies on the Antioxidant Effect of Rutin in the Microenvironment of Cationic Micelles. *Microbiology Acta*, 161: 233-239.
- Jayaprakash, G.K., Selvi, T and Sakariah, K. (2003). Antimicrobial and Antioxidant Activities of Grape (*Vitis Vinifera*) Seed Extracts. *Food Research International*, 36: 117-122.
- Kennedy, D.O and Whightman, E.C. (2011). Herbal Extracts and Phytochemicals: Plant Secondary Metabolites and Enhancement of Human Brain Function. *Advances in Nutrition*, 2: 32-50.
- Kwon, Y.S., Ann, H.S., Nabeshima, T., Shin, E.J., Kim, W.K and Jhoo J.H. (2004). Selegiline Potentiates the Effects of EGb 761 in Response to Ischemic Brain Injury. *Neurochemistry International Journal*, 45: 157-170.
- Lacaille-Dubois, M.A and Wagner, H. (2000). Bioactive Saponin from Plants: An Update. Studies in Natural Product Chemistry, Atta-ur-Rahman ed. *Elsevier Science*, 21: 633-687.
- Latge, J.P. (2007). The Cell Wall: A Carbohydrate Armour for Fungal Cell. *Molecular Microbiology*, 66(2): 279-290.
- Martin, A.J.P and Synge, R.L.M. (1941). A New Form of Chromatogram Employing Two Liquid Phases. *Journal of Biochemistry*, 35: 1358-1560.
- Pallota, R.C, Ribeiro, M.S and De lima Machado, M.E. (2007). Determination of the Minimal Inhibitory Concentration of Four Medicaments Used as Intracanal Medication. *Australian Endodontics Journal*, 33: 107-111.
- Peach, K.C., Bray, W.M., Winslow, D., Linington, P.F and Linington R.G. (2013). Mechanism of Action-Based Classification of Antibiotics Using High-Content Bacterial Image Analysis. *Molecular Biosystems*, 9(7): 1837-1848.
- Price, K.R., Johnson, I.J. and Frenwick, G.R. (1987). The Chemistry and Biological Significance of Saponin in Foods and Feeding Stuffs. *Critical Review of Food Science and Nutrition*, 26: 633-87
- Raaman, N. (2008). Phytochemical Techniques. New India Publishing Agency, New-Delhi, India. Pp 1-9.
- Sati, P., Pandey, A., Rawat, S and Rani, A. (2013). Phytochemicals and Antioxidants in Leaf Extracts of *Ginkgo biloba* with Reference to Location, Seasonal Variation and Solvent System. *Journal of Pharmacy Research*, 7: 804-809
- Sati, P., Pandey, A and Palni, L.M.S. (2012). Antimicrobial Potential of Leaf Extracts of *Ginkgo biloba L.* Growing in Uttarakhand, India. *National Academy of Science Letters*, 35(3): 201-206.
- Sati, S.C and Joshi, S. (2011). Antibacterial Activities of *Ginkgo biloba L.* Extracts. *Scientific World Journal*, 11: 2337-2242.
- Tyagi, E., Agrawal, R., Nath, C and Shukla, R. (2010). Cholinergic Protection via Alpha 7 Nicotinic Acetylcholinergic Receptors and PI3K-AKT Pathway in LPS-induced Neuroinflammation. *Neurochemistry International*, 56: 135-142.

- Ujowundu, F.N., Ukoha, A.I., Ojiako, A.O and Nwaoguikpe, R.N. (2015).Isolation of Bioactive Phytochemicals in Leaves of *Combretum dolichopetalum* and their Hydrogen Peroxide Scavenging Potentials.*Pharmaceutica Analytica Acta Journal*,**6**:435-444.
- Wadsworth, T.I and Koop, D.R. (2001).Effects of *Ginkgo biloba* Extract (EGb 761) and Quercetin on Lipopolysaccharide-Induced Release of Nitric Oxide.*Chemico-Biological Interaction*,**137**: 43-58.